



Epidermal Nerve Fiber Density Testing

Clinical Policy ID: CCP.1263

Recent review date: 6/2026

Next review date: 10/2027

Policy contains: Epidermal nerve fiber density testing; skin punch biopsy; small fiber neuropathy.

Select Health of South Carolina has developed clinical policies to assist with making coverage determinations. Select Health of South Carolina's clinical policies are based on guidelines from established industry sources, such as the Centers for Medicare & Medicaid Services (CMS), state regulatory agencies, the American Medical Association (AMA), medical specialty professional societies, and peer-reviewed professional literature. These clinical policies along with other sources, such as plan benefits and state and federal laws and regulatory requirements, including any state- or plan-specific definition of "medically necessary," and the specific facts of the particular situation are considered, on a case by case basis, by Select Health of South Carolina when making coverage determinations. In the event of conflict between this clinical policy and plan benefits and/or state or federal laws and/or regulatory requirements, the plan benefits and/or state and federal laws and/or regulatory requirements shall control. Select Health of South Carolina's clinical policies are for informational purposes only and not intended as medical advice or to direct treatment. Physicians and other health care providers are solely responsible for the treatment decisions for their patients. Select Health of South Carolina's clinical policies are reflective of evidence-based medicine at the time of review. As medical science evolves, Select Health of South Carolina will update its clinical policies as necessary. Select Health of South Carolina's clinical policies are not guarantees of payment.

Coverage policy

Epidermal nerve fiber density testing by skin biopsy is clinically proven and, therefore, may be medically necessary for the detection of small fiber neuropathy when **all** of the following criteria are met (American Association of Neuromuscular and Electrodiagnostic Medicine [Katzberg, 2026]; England, 2009, reaffirmed 2025):

- Member presents with symptoms of painful sensory neuropathy
- Member has no history of a disorder known to predispose to painful neuropathy (e.g., diabetic neuropathy).
- No evidence of large-fiber neuropathy on **both**:
 - Physical examination (e.g., reduced or absent muscle-stretch reflexes or reduced proprioception and vibration sensation)
 - Electromyography and nerve-conduction studies

Limitations

All other uses of epidermal nerve fiber density testing are investigational/not clinically proven and, therefore, not medically necessary.

Alternative covered services

- Neurologic consultation
- Screening for other treatable causes of small fiber neuropathy
- Functional tests (e.g., quantitative sensory testing)
- Autonomic testing
- Nerve conduction testing
- Somatosensory evoked potentials
- Nerve biopsy

Background

Small fiber neuropathy, also known as small-fiber sensory/peripheral neuropathy, is a peripheral nerve disease that selectively affects small diameter myelinated and non-myelinated nerve fibers (Cascio, 2022). Sensory symptoms of small fiber neuropathy vary widely in pattern and severity. Treatment is generally palliative, and not curative.

Etiologies associated with small fiber neuropathy include genetic mutations in the *SCN9A* or *SCN10A* gene, diabetes, impaired glucose tolerance, several hereditary disorders, certain autoimmune disorders, viral and infectious diseases (e.g., human immunodeficiency virus infection), neurotoxic medications, and alcoholism (Cascio, 2022; Görlach, 2020; Raicher, 2022). In up to 50% of cases, the etiology is idiopathic and often presents as burning feet. With repeated skin biopsies over 12 to 18 months, an intraepidermal decrease in nerve fiber density is seen in the legs, which demonstrates the progressive degeneration of this condition (Raicher, 2022).

Small fiber neuropathy is a diagnosis of exclusion based on clinical findings and the absence of large fiber involvement, particularly in the context of an associated disease, such as diabetes. Ancillary testing and specialty consultation may provide additional guidance. Testing includes screening for other treatable causes of small fiber neuropathy, scoring examinations, and characterizing specific types of pain and genetic testing. Electromyography and nerve conduction studies assess possible larger myelinated sensory and motor fiber involvement (Cascio, 2022).

Epidermal nerve fiber density testing, also called intraepidermal nerve fiber density testing, assesses the structural integrity of small nerve fibers using skin biopsy and immunostaining (Cascio, 2022; Raicher, 2022). It quantifies the intraepidermal nerve fibers crossing the epidermis, and results are expressed as the number of intraepidermal nerve fibers per millimeter. Epidermal nerve fiber density testing is regulated under the Clinical Laboratory Improvement Amendments of 1988 (42 U.S.C. 263a).

Normative values vary depending on sampling site, quantification technique, patient age, and gender. Laboratories may use established normative values or develop their own methods for determining reference ranges and cutoff values. Epidermal nerve fiber density below a normal reference range suggests peripheral neuropathy, raising the suspicion of disorders known to cause small fiber neuropathy such as diabetes, impaired glucose tolerance and certain autoimmune diseases. Epidermal nerve fiber density within the normal range suggests the need to test for etiologies other than those known to produce peripheral neuropathy. In addition, epidermal nerve fiber density testing may be used to assess morphological changes of intraepidermal nerve fibers and dermal nerve fibers (Cascio, 2022; Görlach, 2020).

Findings

Epidermal nerve fiber density with skin punch biopsy using bright-field immunohistochemistry is a safe procedure with no major complications and for which normative data exist to characterize findings as normal or abnormal. Epidermal nerve fiber density testing has a high diagnostic yield¹ (in this case, equivalent to sensitivity) for identifying pathologic changes in unmyelinated small nerve fibers. Presently, the true value of epidermal nerve fiber density for diagnosing sensory neuropathy depends on its ability to distinguish patients with small fiber neuropathy from patients whose symptoms are unrelated to neuropathy. Therefore, there is sufficient evidence to support using epidermal nerve fiber density testing to rule out non-neuropathic involvement in patients with symptoms suggestive of small fiber neuropathy who have no evidence of large fiber neuropathy and no disorder known to predispose to painful neuropathy.

Guidelines

The American Association of Neuromuscular and Electrodiagnostic Medicine Small Fiber Neuropathy Task Force stated that intraepidermal nerve fiber density testing at the distal leg site is probably useful in identifying patients with small fiber neuropathy (Level B), based on two Class II and two Class III studies demonstrating that reduced intraepidermal nerve fiber density at a standard site 10 cm from the lateral malleolus can identify patients with small fiber neuropathy. Test sensitivity ranged from 74% to 78%, and specificity ranged from 65% to 80%. Nerve fiber morphology, nerve fiber length, and interfiber distance are possibly useful in identifying small fiber neuropathy (Level C). There is insufficient evidence to comment on the use of biopsy obtained from proximal thigh sites in the evaluation of small fiber neuropathy (Level U) (Katzberg, 2026).

A joint guideline by the American Academy of Neurology, American Association of Neuromuscular and Electrodiagnostic Medicine, and American Academy of Physical Medicine and Rehabilitation further examined the ability of intraepidermal nerve fiber density to distinguish symptomatic patients with polyneuropathy from symptomatic patients without polyneuropathy. In this setting, the sensitivity for the diagnosis of polyneuropathy ranged from 0.45 to 0.90, but the specificity was consistently higher, ranging from 0.95 to 0.97, raising the likelihood of polyneuropathy in the presence of reduced intraepidermal nerve fiber density. Therefore, skin biopsy may be considered in symptomatic patients with suspected polyneuropathy to diagnose the presence of a polyneuropathy, particularly small fiber sensory polyneuropathy (England, 2009, reaffirmed 2025).

Evidence review

We identified six individual studies (Caro, 2014; Grone, 2014; Kim, 2014; Kosmidis, 2014; Shikuma, 2015; Timar, 2016) for this policy. The best available evidence for epidermal nerve fiber density testing consists of case-control and cross-sectional studies of patients with clinical sensory neuropathy referred to neurology specialty clinics compared to healthy controls. The remaining studies were of insufficient quality and quantity to assess the ability of epidermal nerve fiber density testing to detect preclinical neuropathy in persons with known disease and mixed neuropathy status, disease severity, or response to treatment. No studies have assessed the ability of epidermal nerve fiber density testing to distinguish disease etiology, change clinical management (particularly in the presence of known causes of neuropathy such as diabetes), or improve patient outcomes.

One systematic review examining the diagnostic criteria for idiopathic small fiber neuropathy highlighted the need to develop standardized, evidence-based guidelines (Haroutounian, 2021).

¹ I.e., the probability that epidermal nerve fiber density will be abnormal in a particular population. A high diagnostic yield would limit the number of patients in whom underlying causes other than peripheral neuropathy need to be investigated. It may or may not provide useful prognostic information beyond that obtained from basic clinical measurements.

Løseth (2024) analyzed the extent to which epidermal nerve fiber density testing and quantitative sensory testing were abnormal in an unselected cohort (n = 203) of participants with symptoms suggestive of small fiber neuropathy and normal nerve conduction studies. The most prevalent underlying conditions were diabetes mellitus, cancer/cytostatics, sarcoidosis, fibromyalgia, hypothyreosis, and Sjögren syndrome, but 113 (55.7%) participants had no established cause. Less than half (45.3 %) had reduced epidermal nerve fiber density, and 50% had abnormal quantitative sensory testing. There were no gender differences in epidermal nerve fiber density testing results.

In 2017, we identified no new information for the policy, and no policy changes are warranted.

In 2018, we identified no new information to add to the policy, and no policy changes are warranted. The policy ID was changed from CP# 09.01.12 to CCP.1263.

In 2019, we identified no new relevant information to add to the policy. No policy changes are warranted.

In 2020, we identified no new relevant information to add to the policy. No policy changes are warranted.

In 2021, we updated the references with no policy changes warranted.

In 2022, we identified no new relevant information to add to the policy. No policy changes are warranted.

In 2024, we updated the references and made no policy changes.

In 2025, we updated the references and identified no new relevant information to add to the policy. No policy changes are warranted.

In 2026, we updated the references and added one new guideline to the policy. No policy changes are warranted

References

On May 12, 2026, we searched PubMed and the databases of the Cochrane Library, the U.K. National Health Services Centre for Reviews and Dissemination, the Agency for Healthcare Research and Quality, and the Centers for Medicare & Medicaid Services. Search terms were "Nerve Fibers (MeSH)," "Epidermis (MeSH)," "small fiber neuropathy (MeSH)," and the free text term "epidermal nerve fiber density." We included the best available evidence according to established evidence hierarchies (typically systematic reviews, meta-analyses, and full economic analyses, where available) and professional guidelines based on such evidence and clinical expertise.

42 U.S.C. 263a. Certification of laboratories.

Caro XJ, Winter EF. Evidence of abnormal epidermal nerve fiber density in fibromyalgia: Clinical and immunologic implications. *Arthritis Rheumatol*. 2014;66(7):1945-1954. Doi: 10.1002/art.38662.

Cascio MA, Mukhdomi T. Small Fiber Neuropathy. In: *StatPearls* [Internet]. Treasure Island (FL): StatPearls Publishing; 2026 Jan-. <https://www.ncbi.nlm.nih.gov/books/NBK582147/>. Updated December 12, 2022.

England JD, Gronseth GS, Franklin G, et al. Practice parameter: Evaluation of distal symmetric polyneuropathy: Role of autonomic testing, nerve biopsy, and skin biopsy (an evidence-based review). Report of the American Academy of Neurology, American Association of Neuromuscular and Electrodiagnostic Medicine, and American Academy of Physical Medicine and Rehabilitation. *Neurology*. 2009;72(2):177-184. Doi: 10.1212/01.wnl.0000336345.70511.0f. <https://www.aan.com/Guidelines/home/GuidelineDetail/315>. Reaffirmed February 8, 2025.

Görlach J, Amsel D, Kölbel H, et al. Diagnostic utility of small fiber analysis in skin biopsies from children with chronic pain. *Muscle Nerve*. 2020;61(2):173-181. Doi: 10.1002/mus.26766.

- Grone E, Uceyler N, Abahji T, et al. Reduced intraepidermal nerve fiber density in patients with chronic ischemic pain in peripheral arterial disease. *Pain*. 2014;155(9):1784-1792. Doi: 10.1016/j.pain.2014.06.003.
- Haroutounian S, Todorovic MS, Leinders M, et al. Diagnostic criteria for idiopathic small fiber neuropathy: A systematic review. *Muscle Nerve*. 2021;63(2):170-177. Doi: 10.1002/mus.27070.
- Katzberg HD, So Y, Brannagan T, et al. Diagnostic and screening laboratory tests in the assessment of patients with small fiber neuropathy: An evidence-based review-report of the American Association of Neuromuscular and Electrodiagnostic Medicine Small Fiber Neuropathy Task Force. *Muscle Nerve*. 2026;73(6):952-960. Doi: 10.1002/mus.70119.
- Kim TW, Shim WH, Kim JM, et al. Clinical characteristics of pruritus in patients with scalp psoriasis and their relation with intraepidermal nerve fiber density. *Ann Dermatol*. 2014;26(6):727-732. Doi: 10.5021/ad.2014.26.6.727.
- Kosmidis ML, Koutsogeorgopoulou L, Alexopoulos H, et al. Reduction of intraepidermal nerve fiber density (iENFD) in the skin biopsies of patients with fibromyalgia: A controlled study. *J Neurol Sci*. 2014;347(1-2):143-147. Doi: 10.1016/j.jns.2014.09.035.
- Løseth S, Nebuchennykh M, Brokstad RT, Lindal S, Mellgren SI. Cutaneous nerve biopsy in patients with symptoms of small fiber neuropathy: A retrospective study. *Scand J Pain*. 2024;24(1). Doi: 10.1515/sjpain-2023-0071.
- Raicher I, Casartelli-Ravagnani LH, Correa SG, et al. Investigation of nerve fibers in the skin by biopsy: technical aspects, indications, and contribution to diagnosis of small-fiber neuropathy. *Einstein (Sao Paulo)*. 2022;20:eMD8044. Doi: 10.31744/einstein_journal/2022MD8044.
- Shikuma CM, Bennett K, Ananworanich J, et al. Distal leg epidermal nerve fiber density as a surrogate marker of human immunodeficiency-associated sensory neuropathy risk: Risk factors and change following initial antiretroviral therapy. *J Neurovirol*. 2015;21(5):525-534. Doi: 10.1002/mus.23834.
- Timar B, Popescu S, Timar R, et al. The usefulness of quantifying intraepidermal nerve fibers density in the diagnostic of diabetic peripheral neuropathy: A cross-sectional study. *Diabetol Metab Syndr*. 2016;8;31. Doi: 10.1186/s13098-016-0146-4.

Policy updates

9/2016: initial review date and clinical policy effective date: 1/2017

10/2017: Policy references updated.

10/2018: Policy references updated. Policy ID changed.

10/2019: Policy references updated.

12/2020: Policy references updated.

12/2021: Policy references updated.

12/2022: Policy references updated.

6/2024: Policy references updated.

6/2025: Policy references updated.

6/2026: Policy references updated

Related codes

Below are the most commonly submitted codes for the service(s)/item(s) subject to this policy CCP.1263. This is not an exhaustive list of codes. Providers are expected to consult the appropriate coding manuals and bill accordingly.

Code	Code description
88356	Morphometric analysis; nerve